arXiv:1301.3642v1 [cond-mat.quant-gas] 16 Jan 2013

First-Order Quantum Phase Transition in Bose Gases

Nguyen Thanh Phuc,[1] Yuki Kawaguchi,[2] and Masahito Ueda[1]
[1]*Department of Physics, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*
[2]*Department of Applied Physics and Quantum-Phase Electronics Center, University of Tokyo, 2-11-6 Yayoi, Bunkyo-ku, Tokyo 113-0032, Japan*

(Dated: January 17, 2013)

The Bogoliubov theory has proven to be an accurate and versatile tool in the study of weakly-interacting dilute Bose gases at low temperatures, yet there is one exception where it goes qualitatively wrong, i.e., near the first-order quantum phase transitions. By examining a phase transition in spinor Bose-Einstein condensates (BECs), we find that the energy spectrum given by the Bogoliubov theory is inconsistent with the fact that the phase transition is first order. We resolve this problem by calculating the spectrum based on the spinor version of the Beliaev theory. We also discuss the ground-state phase diagram of spin-2 BECs which is modified by quantum fluctuations and the possibility of macroscopic quantum tunneling near the cyclic-nematic phase boundary.

PACS numbers: 03.75.Mn,03.75.Kk,67.85.Jk

Quantum phase transitions have been an active field of research [1, 2], and played a central role in the study of electrical and magnetic properties of solid-state materials such as magnetic insulators, heavy fermion materials, novel semiconductor structures, and high-temperature superconductors. In ultracold atoms, the quantum phase transition from superfluids to Mott insulators has been extensively investigated both theoretically and experimentally [3, 4]. Although the focus of these studies has been mainly on the second-order or continuous quantum phase transitions, the first-order quantum phase transitions in fermionic systems such as itinerant-electron magnets [5] and superfluid helium 3 [6] have recently attracted much attention in connection with an extended non-Fermi liquid phase and superconductivity [7, 8].

In this Letter, we point out that the Bogoliubov theory [9], which has been widely used to investigate weakly-interacting dilute Bose gases [10], is inadequate to describe spinor Bose-Einstein condensates (BECs). In spinor BECs, there exist several ground-state phases with distinct symmetries [11]. The phase transitions between them, therefore, should be first order. However, the result of a Bogoliubov analysis implies that there is no metastable state around the phase boundary, suggesting that the phase transition is second order. We resolve this inconsistency by invoking the spinor version of the Beliaev theory [12, 13]. We will elucidate the above statement by taking spin-2 Bose gases as an example. After showing that quantum fluctuations modify the mean-field phase diagram of spin-2 BECs, we examine in detail the transition between the uniaxial nematic and cyclic phases and prove that the Beliaev energy spectrum actually supports metastable states. We argue that this should be a generic feature of first-order quantum phase transitions in bosonic systems since the energy landscape of a first-order quantum phase transition requires higher-order terms that are beyond the ϕ^4 model and Bogoliubov theory. Finally, the possibility of quantum macroscopic tunneling (MQT) from a metastable state to the ground state in the first-order quantum phase transition is discussed.

The Hamiltonian of a homogeneous spin-2 spinor BEC is given by $\hat{H} = \hat{h}_0 + \hat{V}$ with kinetic energy $\hat{h}_0 = \int d\mathbf{r} \sum_{j=-2}^{2} \hat{\psi}_j^\dagger(\mathbf{r})(-\hbar^2\nabla^2/2M)\hat{\psi}_j(\mathbf{r})$ (M is the atomic mass) and interaction energy [14]

$$\hat{V} = \frac{1}{2}\int d\mathbf{r}\Big[c_0 : \hat{n}^2 : + c_1 : \hat{\mathbf{F}}^2 : + c_2 : \hat{A}_{00}^\dagger \hat{A}_{00} :\Big]. \quad (1)$$

Here, :: denotes normal ordering of operators, and $\hat{n} = \sum_j \hat{\psi}_j^\dagger(\mathbf{r})\hat{\psi}_j(\mathbf{r})$, $\hat{\mathbf{F}} = \sum_{i,j} \hat{\psi}_i^\dagger(\mathbf{r})(\mathbf{f})_{ij}\hat{\psi}_j(\mathbf{r})$, and $\hat{A}_{00} = (1/\sqrt{5})\sum_j(-1)^{-j}\hat{\psi}_j(\mathbf{r})\hat{\psi}_{-j}(\mathbf{r})$ are the number density, spin density, and spin-singlet pair amplitude operators, respectively, with $(\mathbf{f})_{ij}$ being the spin-2 matrix vector. The coefficients c_0, c_1, c_2 are related to the s-wave scattering lengths $a_{\mathcal{F}}$ ($\mathcal{F} = 0, 2, 4$) of the total spin $\mathcal{F}$ channel by $c_0 = 4\pi\hbar^2(4a_2 + 3a_4)/(7M)$, $c_1 = 4\pi\hbar^2(a_4 - a_2)/(7M)$, $c_2 = 4\pi\hbar^2(7a_0 - 10a_2 + 3a_4)/(7M)$, respectively. The order parameter is represented by a five-component spinor $\vec{\phi} = \sqrt{n_0}(\xi_2, \xi_1, \xi_0, \xi_{-1}, \xi_{-2})^{\mathrm{T}}$, where n_0 is the number density of condensed atoms and T denotes transpose. At the mean-field level, the ground-state phase diagram obtained by minimizing the energy functional is shown in Fig. 1(a) [14]. There is an energy degeneracy in the manifold of nematic phases with order parameters $\vec{\phi} = \sqrt{n_0}(\sin\eta/\sqrt{2}, 0, \cos\eta, 0, \sin\eta/\sqrt{2})^{\mathrm{T}}$, where η is the degeneracy parameter.

Since the order parameters of different phases cannot be transformed continuously to each other, the phase transitions between them must be first order. Nevertheless, the energy spectrum given by the Bogoliubov theory shows either a dynamical or a Landau instability at all phase boundaries of spinor BECs under zero magnetic field. For example, one of the elementary excitations of the nematic phase has the energy spectrum given by $\hbar\omega_{\mathbf{p}} = \sqrt{\epsilon_{\mathbf{p}}^0(\epsilon_{\mathbf{p}}^0 - 2c_2 n)}$, where $\epsilon_{\mathbf{p}}^0 \equiv \hbar^2\mathbf{p}^2/(2M)$ and n is the total number density. This spectrum becomes a complex number at long wavelengths for positive c_2, indicating a dynamical instability of the nematic phase. Such an instability at the phase boundary, however, characterizes second-order as opposed to first-order phase transitions with the existence of metastable states [15]. The Bogoliubov energy spectrum is, therefore, inconsistent with the fact that the phase transition is first order.



FIG. 1: Ground-state phase diagram of spin-2 BECs obtained with (a) the mean-field approximation and (b) the Lee-Huang-Yang (LHY) correction. The dashed lines in (b) indicate the mean-field phase boundaries in (a). The inset in each phase shows the surface plot of the spinor order parameter defined as $|\psi(\theta,\phi)|^2 \equiv |\sum_m \xi_m Y_m^2(\theta,\phi)|^2$, where the hue indicates the phase of $\psi(\theta,\phi)$ according to the color gauge shown on the right. The LHY corrections due to quantum fluctuations lift the degeneracy in the nematic phase in (a), leading to the uniaxial nematic (UN) and biaxial nematic (BN) phases for $c_1 > 0$ and $c_1 < 0$, respectively. Quantum fluctuations also shift the cyclic-UN and ferromagnetic-BN phase boundaries. However, the cyclic-ferromagnetic phase boundary is not affected by quantum fluctuations because of the destructive interference caused by time-reversal symmetry breaking (see text).

We first show that the mean-field phase diagram is modified due to the Lee-Huang-Yang (LHY) correction, which arises from quantum fluctuations. As shown in Refs. [16, 17], quantum fluctuations lift the energy degeneracy in the manifold of nematic phases, making the uniaxial nematic (UN) phase ($\eta = 0$) and biaxial nematic (BN) phase ($\eta = \pi/2$) the ground states for $c_1 > 0$ and $c_1 < 0$, respectively. Quantum fluctuations also shift the UN-cyclic and BN-ferromagnetic phase boundaries as follows. Up to the LHY correction, the ground-state energy densities of the cyclic and UN phases for $c_1 > 0$, $c_2 \leq 0$ are given by [18]

$$\frac{E^{\mathrm{C}}}{V} = \frac{c_0 n^2}{2} + \frac{8M^{3/2}}{15\pi^2\hbar^3}\left[(nc_0)^{5/2} + 12\sqrt{2}(nc_1)^{5/2}\right], \quad (2)$$

$$\begin{aligned}\frac{E^{\mathrm{UN}}}{V} =& \left(c_0 + \frac{c_2}{5}\right)\frac{n^2}{2} + \frac{8M^{3/2}}{15\pi^2\hbar^3}\left[(nc_0)^{5/2} + 18\sqrt{3}(nc_1)^{5/2}\right] \\ &+ \mathcal{O}\left[\frac{M^{3/2}n^{5/2}\max\left\{c_0^{3/2}, c_1^{3/2}\right\}|c_2|}{\hbar^3}\right], \end{aligned} \quad (3)$$

respectively. Here, we expanded E^{UN} in powers of c_2/c_0 and c_2/c_1, which are expected to be small near the UN-cyclic phase boundary. In fact, since $|c_2| \sim M^{3/2}n^{1/2}c_1^{5/2}/\hbar^3$ at the phase boundary [see Eq. (4)], the last term in Eq. (3) is smaller than the others by a factor of $\sqrt{na^3} \ll 1$ with $a \equiv (4a_2 + 3a_4)/7 = c_0 M/(4\pi\hbar^2)$, and thus can be neglected. By comparing the energies in Eqs. (2) and (3), we find that the phase boundary between the UN and cyclic phases is shifted to

$$\begin{aligned} c_2^{\mathrm{UN-C}} &\simeq -\frac{16(18\sqrt{3} - 12\sqrt{2})M^{3/2}n^{1/2}c_1^{5/2}}{3\pi^2\hbar^3} \\ &\simeq -342\left(\frac{c_1}{c_0}\right)^{3/2}\sqrt{na^3}\, c_1, \end{aligned} \quad (4)$$

i.e., the region of the cyclic phase is expanded. Similarly, the ground-state energies of the ferromagnetic and BN phases with LHY corrections for $c_1 < 0$, $c_2 \simeq 20c_1$ are given by [18]

$$\frac{E^{\mathrm{F}}}{V} = \left(\frac{c_0}{2} + 2c_1\right)n^2\left[1 + \frac{16M^{3/2}}{15\pi^2\hbar^3}\sqrt{n(c_0+4c_1)^3}\right], \quad (5)$$

$$\begin{aligned}\frac{E^{\mathrm{BN}}}{V} =& \left(\frac{c_0}{2} + \frac{c_2}{10}\right)n^2\left[1 + \frac{16M^{3/2}}{15\pi^2\hbar^3}\sqrt{n\,(c_0+4c_1)^3}\right] \\ &+ \frac{8M^{3/2}}{15\pi^2\hbar^3}(32 + 18\sqrt{3})(|c_1|n)^{5/2} \\ &+ \mathcal{O}\left[\frac{M^{3/2}n^{5/2}\max\left\{c_0^{3/2}, |c_1|^{3/2}\right\}|c_2 - 20c_1|}{\hbar^3}\right], \end{aligned} \quad (6)$$

respectively. By noting that $|c_2 - 20c_1| \sim M^{3/2}n^{1/2}|c_1|^{5/2}/\hbar^3$ near the phase boundary [Eq. (7)], the last term in Eq. (6) is smaller than the others by a factor of $\sqrt{na^3} \ll 1$, and thus negligible. Consequently, the phase boundary between the ferromagnetic and BN phases is shifted from its mean-field value $c_2 = 20c_1$ to

$$\begin{aligned} c_2^{\mathrm{F-BN}} &\simeq 20c_1 - \frac{32(16+9\sqrt{3})M^{3/2}n^{1/2}|c_1|^{5/2}}{3\pi^2\hbar^3} \\ &\simeq 20c_1 - 1521\left(\frac{c_1}{c_0}\right)^{3/2}\sqrt{na^3}\,c_1, \end{aligned} \tag{7}$$

i.e., the region of the ferromagnetic phase is expanded.

On the other hand, the phase boundary between the ferromagnetic and cyclic phases at $c_1 = 0$ and $c_2 > 0$ is not influenced by quantum fluctuations since both of the ground-state energies are independent of c_2, and equal to each other at $c_1 = 0$ [see Eqs. (2) and (5)]. That E^{C} and E^{F} are independent of c_2 is due to time-reversal symmetry breaking in these phases. For the ferromagnetic phase with order parameter $\vec{\phi} = \sqrt{n_0}(1,0,0,0,0)^{\mathrm{T}}$, excitations caused by interaction c_2 such as $(\mathbf{p}=\mathbf{0}, m_F=2)+(\mathbf{p}=\mathbf{0}, m_F=-2) \to (\mathbf{p}, m_F=1)+(-\mathbf{p}, m_F=-1)$ do not occur due to the absence of condensed atoms in the magnetic sublevel $m_F = -2$. Meanwhile, the cyclic phase has the order parameter $\vec{\phi} = \sqrt{n_0}(1/2, 0, \sqrt{2}/2, 0, -1/2)^{\mathrm{T}}$ with opposite sign for $m_F = \pm 2$ components, which results from time-reversal symmetry breaking. Therefore, the amplitudes of excitations from $(\mathbf{p}=\mathbf{0}, m_F=2)+(\mathbf{p}=\mathbf{0}, m_F=-2)$ and from $(\mathbf{p}=\mathbf{0}, m_F=0)+(\mathbf{p}=\mathbf{0}, m_F=0)$ to $(\mathbf{p}, m_F=1)+(-\mathbf{p}, m_F=-1)$ both via interaction c_2 interfere destructively with each other.

In the presence of an external magnetic field, the difference in the LHY correction among ground-state phases, whose magnitude is of the order of $\Delta E \equiv M^{3/2}c_1^{5/2}n^{3/2}/\pi^2\hbar^3$, can compete with the quadratic Zeeman energy q_B, and the phase diagram depends on the relative strength of these two effects. In the limit of a high magnetic field $q_B \gg \Delta E$, the effect of quantum fluctuations can be neglected, and the ground-state phase diagram is obtained by mean-field theory [19]. In the opposite limit of low magnetic field $q_B \ll \Delta E$, the quantum fluctuations dominate, the quadratic Zeeman energy becomes negligible, and the ground-state phase diagram would be Fig. 1(b). These two distinct regimes can be crossed over at $q_B \sim \Delta E$, which corresponds to a magnetic field of the order of 7 mG by using the parameters of ^{87}Rb [20, 21] and atomic density $n = 10^{15}\,\mathrm{cm}^{-3}$. If an external magnetic field is made as small as 0.1 mG [22], all the above regimes can be investigated experimentally.

The inconsistency between the energy spectrum given by the Bogoliubov theory and the phase diagram even becomes more serious with the addition of LHY correction to the ground-state energy. Indeed, the cyclic phase is the ground state for $c_2^{\mathrm{UN-C}} < c_2 < 0$ whereas the Bogoliubov energy spectrum shows an instability of the cyclic phase for $c_2 < 0$. Now we show that all the above inconsistencies are resolved if we go to the next level of approximation, i.e., with the Beliaev theory [12, 13]. The Green's function-based Beliaev theory is applied to calculate the second-order energy spectrum of the UN phase in a manner similar to our previous work on spin-1 spinor BECs [23]. The Dyson's equation for the Green's functions is given by

$$G_{jj'}^{\alpha\beta}(p) = (G^0)_{jj'}^{\alpha\beta}(p) + (G^0)_{jm}^{\alpha\gamma}\Sigma_{mm'}^{\gamma\delta}(p)G_{m'j'}^{\delta\beta}(p), \tag{8}$$

where the interacting Green's functions, non-interacting Green's functions, and self-energies are denoted by G, G^0, and Σ, respectively, all of which are 10×10 matrices with $j, j', m, m' = -2, \dots, 2$ indicating the spin index and $\alpha, \beta, \gamma, \delta = 1, 2$ accounting for the normal and anomalous components. For the UN phase, each of the 10×10 matrices reduces to five 2×2 submatrices, which facilitates the analytic calculation. The zero-momentum energy of the elementary excitation that drives the transition from the UN to cyclic phase is then calculated to be

$$\hbar\omega_{\mathbf{p}=\mathbf{0}} \simeq \frac{\sqrt{\left[-\frac{c_2 n_0}{5} + A - \mu^{(2)}\right]^2 - \left[\frac{c_2 n_0}{5} + C\right]^2}}{1-B}, \tag{9}$$

where $\mu^{(2)}$ is the second-order chemical potential, and A, B, and C are the lowest-order coefficients in the frequency-power expansion of the second-order self-energies $\Sigma_{2,2}^{11(2)}(p)$, $\Sigma_{2,2}^{22(2)}(p)$, and $\Sigma_{2,-2}^{12(2)}(p)$, respectively. These coefficients are given by

$$\frac{A-\mu^{(2)}}{(Mn_0)^{3/2}} \simeq -\frac{4\sqrt{3}c_1^{5/2}}{\pi^2\hbar^3} + \frac{\left(42\sqrt{3}c_1^{3/2} - 10c_0^{3/2}\right)c_2}{15\pi^2\hbar^3}, \tag{10}$$

$$\frac{C}{(Mn_0)^{3/2}} \simeq \frac{12\sqrt{3}c_1^{5/2}}{\pi^2\hbar^3} + \frac{\left(10c_0^{3/2} - 30\sqrt{3}c_1^{3/2}\right)c_2}{15\pi^2\hbar^3}, \tag{11}$$

$$\frac{B}{M^{3/2}n_0^{1/2}} \simeq -\frac{\left(c_0^{3/2} + 3\sqrt{3}c_1^{3/2}\right)}{3\pi^2\hbar^3} - \frac{\left(c_0^{1/2} + \sqrt{3}c_1^{1/2}\right)c_2}{30\pi^2\hbar^3}. \tag{12}$$

Here, using the same argument as below Eq. (3), all terms with higher powers of c_2 are smaller than those in Eqs. (10)-(12) by at least a factor of $\sqrt{na^3} \ll 1$, and thus neglected. By substituting Eqs. (10)-(12) into Eq. (9), we find that the UN phase becomes dynamically unstable if

$$\begin{aligned} c_2 > c_2^{\mathrm{UN-unstable}} &\equiv -\frac{40\sqrt{3}M^{3/2}n^{1/2}c_1^{5/2}}{\pi^2\hbar^3} \\ &\simeq -313\left(\frac{c_1}{c_0}\right)^{3/2}\sqrt{na^3}\,c_1. \end{aligned} \tag{13}$$

From Eqs. (4) and (13), we see that the UN phase becomes metastable for

$$-342\left(\frac{c_1}{c_0}\right)^{3/2}\sqrt{na^3}\,c_1 < c_2 < -313\left(\frac{c_1}{c_0}\right)^{3/2}\sqrt{na^3}\,c_1. \tag{14}$$

The existence of this metastable state and the associated hysteresis phenomenon are characteristic of first-order phase transitions [15]. Similarly, a metastable state of the cyclic phase is expected to exist in the parameter regime $c_2^{\mathrm{UN-C}} > c_2 > c_2^{\mathrm{C-unstable}}$. Consequently, the inconsistency of the Bogoliubov energy spectrum with the first-order phase transition and the ground-state phase diagram has been resolved by using the Beliaev theory.

We notice that such a disappearance of metastable states should be a generic feature of the Bogoliubov theory for any bosonic system. The Bogoliubov theory is included by the dynamical form of the mean-field theory characterized by the time-dependent Gross-Pitaevskii equation [10]. At this level of approximation, the field operator is replaced by the order parameter, i.e., the condensate wave function ϕ, and the energy functional is described by a ϕ^4 model. On the other hand, in the case of a first-order phase transition, terms beyond ϕ^4 become relevant [15], and they can only be obtained by using models such as the Beliaev theory that take into account quantum fluctuations.

To make a quantitative comparison with other first-order phase transitions, we now evaluate the size of the first-order quantum phase transition between UN and cyclic phases. For a first-order thermal phase transition, its size is defined by the ratio of the latent heat to the jump in the specific heat and shown to be proportional to the size of the temperature regime of the metastable state, i.e., the difference between the transition temperature T^{c} and the temperature T^{unstable} at which the instability occurs [24]. Renormalized by T^{c}, the dimensionless size $\delta \equiv |(T^{\mathrm{c}} - T^{\mathrm{unstable}})/T^{\mathrm{c}}|$ for the normal to superconducting transition is $\delta \sim 10^{-6}$, and that for the nematic to smectic-A phase transition in liquid crystals is $\delta \sim 10^{-4}$. This indicates that they are weak first-order phase transitions induced by thermal fluctuations [24]. Similarly, the dimensionless size of the first-order quantum phase transition under consideration can be evaluated as

$$\delta \equiv \left| \frac{c_2^{\mathrm{UN-C}} - c_2^{\mathrm{UN-unstable}}}{c_2^{\mathrm{UN-C}}} \right|. \tag{15}$$

Using Eqs. (4) and (13), we obtain $\delta \simeq 0.1$. This value is much larger than the corresponding values of δ for the above thermal phase transitions, indicating that the quantum phase transition between the UN and cyclic phases with distinct order parameters is a rather strong first-order phase transition.

From Eq. (14), there is a parameter regime in which the UN phase is metastable while the cyclic phase is the ground state. By neglecting a small quantum depletion, these states are approximately given by

$$|\mathrm{UN}\rangle \simeq \left(\hat{a}_0^\dagger\right)^N |\mathrm{vac}\rangle, \tag{16a}$$

$$|\mathrm{Cyclic}\rangle \simeq \left(\frac{\hat{a}_2^\dagger}{2} + \frac{\hat{a}_0^\dagger}{\sqrt{2}} + \frac{\hat{a}_{-2}^\dagger}{2}\right)^N |\mathrm{vac}\rangle, \tag{16b}$$

where $\hat{a}_m^\dagger$ is the creation operator of a particle with zero momentum and magnetic quantum number $m_F = m$ for the vacuum state $|\mathrm{vac}\rangle$. Because these states are not exact eigenstates of the many-body Hamiltonian (1), they can undergo quantum diffusions in the spin space [25–28], and induce a MQT. We now estimate the time scale of the MQT by using a two-state model, i.e., by restricting the Hilbert space to these two states with local energy minima. The time scale of the MQT is then given by $\tau = \hbar/\Delta$ with $\Delta = 2\langle \mathrm{Cyclic}|\hat{V}|\mathrm{UN}\rangle$. Using Eqs. (16) and (1), we obtain

$$\tau \simeq \hbar \frac{2^{N/2}}{c_0 n (N-1)}, \tag{17}$$

where N is the total number of particles. The exponentially large factor of $2^{N/2}$ reflects the macroscopic magnification in a Bose-Einstein condensate. To observe the MQT, τ must be equal to or smaller than the lifetime of the BEC, which is of the order of a second. Substituting parameters of ^{87}Rb into Eq. (17), we obtain the upper limit for the total number of particles $N_{\max} \simeq 36$ for $\tau \lesssim 1$ s.

In conclusion, by examining spin-2 BECs we have shown that the Bogoliubov theory is, in general, inadequate in identifying the nature and boundaries of quantum phase transitions and we should invoke the Beliaev theory to resolve this problem. The ground-state phase diagram is shown to be significantly affected by the quantum interference of virtual excitations and time-reversal symmetry breaking. This study has shed light on the pivotal role of quantum fluctuations in the first-order quantum phase transitions. It will be an interesting future problem to investigate the implication of this study to closely related problems such as the Coleman-Weinberg mechanism of quantum symmetry breaking [29], quantum anomaly [30], and quasi-Nambu-Goldstone modes [31].

This work was supported by KAKENHI (22340114, 22740265, 22103005), a Global COE Program “the Physical Sciences Frontier”, and the Photon Frontier Network Program, from MEXT of Japan. YK acknowledges the financial support from Inoue Foundation.

[1] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, New York, 2011), 2nd edition.

[2] L. D. Carr (ed.), *Understanding Quantum Phase Transitions* (CRC Press, 2010).

[3] M. Greiner, O Mandel, T. Esslinger, T. W. Hansch, and I. Bloch, Nature **415**, 39 (2002)

[4] I. Bloch, J. Dalibard, and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2006).

[5] C. Pfleiderer, J. Phys.: Condens. Matter **17**, S987-S997 (2005).

[6] D. Vollhardt and P. Woelfle, *The Superfluid Phases of Helium 3* (Taylor and Francis, London, 2002).

[7] N. Doiron-Leyraud, I. R. Walker, L. Taillefer, M. J. Steiner, S. R. Julian, and G. G. Lonzarich, Nature **425**, 595 (2003).

[8] S. S. Saxena *et al.*, Nature **406**, 587 (2000).

[9] N. Bogoliubov, J. Phys. USSR 11 (1947) 23.

[10] C. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Bose Gases* (Cambridge University Press, New York, 2008), 2nd edition.

[11] Y. Kawaguchi and M. Ueda, Phys. Rep. **520**, 253 (2012).

[12] S. T. Beliaev, Soviet Physics JETP **7**, 299 (1958).

[13] S. T. Beliaev, Soviet Physics JETP **7**, 289 (1958).

[14] C. V. Ciobanu, S. K. Yip, and T. L. Ho, Phys. Rev. A **61**, 033607 (2000).

[15] K. Huang, *Statistical Mechanics* (John Wiley and Sons, Inc., New York, 1987), 2nd edition.

[16] J. L. Song, G. W. Semenoff, and F. Zhou, Phys. Rev. Lett. **98**, 160408 (2007).

[17] A. M. Turner, R. Barnett, E. Demler, and A. Vishwanath, Phys. Rev. Lett. **98**, 190404 (2007).

[18] S. Uchino, M. Kobayashi, and M. Ueda, Phys. Rev. A **81**, 063632 (2010).

[19] H. Saito and M. Ueda, Phys. Rev. A **72** 053628 (2005).

[20] A. Widera, F. Gerbier, S. Folling, T. Gericke, O. Mandel, and I. Bloch, New J. Phys. **8**, 152 (2006).

[21] E. G. M. van Kempen, S. J. J. M. F. Kokkelmans, D. J. Heinzen, and B. J. Verhaar, Phys. Rev. Lett. **88**, 093201 (2002).

[22] B. Pasquiou, E. Marechal, L. Vernac, O. Gorceix, and B. Laburthe-Tolra, Phys. Rev. Lett. **108**, 045307 (2012).

[23] N. T. Phuc, Y. Kawaguchi and M. Ueda, Ann. Phys. **328**, 158 (2013).

[24] B. I. Halperin, T. C. Lubensky, and S. -K. Ma , Phys. Rev. Lett. **32**, 292 (1974).

[25] C. K. Law, H. Pu, and N. P. Bigelow, Phys. Rev. Lett. **81**, 5257 (1998).

[26] H. Pu, C. K. Law, S. Raghavan, J. H. Eberly, and N. P. Bigelow, Phys. Rev. A **60**, 1463 (1999).

[27] X. Cui, Y. Wang, and F. Zhou, Phys. Rev. A **78** 050701 (2008).

[28] R. Barnett, H. -Y. Hui, C. -H. Lin, J. D. Sau, and S. DasSarma, Phys. Rev. A **83**, 023613 (2011).

[29] S. Coleman and E. Weinberg, Phys. Rev. D **7**, 1888 (1973).

[30] M. Olshanii, H. Perrin, and V. Lorent, Phys. Rev. Lett. **105**, 095302 (2010).

[31] S. Uchino, M. Kobayashi, M. Nitta, and M. Ueda, Phys. Rev. Lett. **105**, 230406 (2010).